



Catalog: OM296274

Scan to get more validated information



Anti-mouse IgG (H+L), F(ab')₂ Fragment (Alexa Fluor® 555 Conjugate)

Catalog: OM296274

☐ 100ug

Product profile

Product name	Anti-mouse IgG (H+L), F(ab') ₂ Fragment (Alexa Fluor® 555 Conjugate)
Antibody Type	Secondary Antibodies
Product description	Anti-Mouse IgG (H+L) F(ab') ₂ Fragment was conjugated to Alexa Fluor® 555 fluorescent dye under optimal conditions and formulated at 2 mg/ml. This F(ab') ₂ fragment results in less non-specific binding to cells through Fc receptors.


Key Feature

Clonality	Monoclonal
Tested Applications	WB ,IP ,IHC ,IF ,FC ,ChIP
Species Reactivity	Mouse
Concentration	1 mg/ml
Purification	

Target Information

Tissue Specificity	F(ab') ₂ fragments are prepared from goat antibodies that have been adsorbed against human IgG and human serum.
--------------------	--

Application



Application

Confocal immunofluorescent analysis of HeLa cells using S6 Ribosomal Protein (54D2) Mouse mAb detected with Anti-Mouse IgG (H+L), F(ab')₂ Fragment (Alexa Fluor® 555 Conjugate) (red, left) compared to an isotype control (right). Actin filaments have been labeled with fluorescein phalloidin. Blue pseudocolor = DRAQ5® (fluorescent DNA dye).

Additional Information

Storage Instructions

The optimal dilution of the anti-species antibody should be determined for each primary antibody by titration. However, a final dilution of 1:250 - 1:1000 should yield acceptable results for immunofluorescent assays.

Storage: Supplied in 0.1 M sodium phosphate, 0.1 M sodium chloride, pH 7.5, 5 mM sodium azide. Store at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.

Storage Buffer

phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Note

The product is for research use only, not for use in diagnostic or therapeutic procedures.

OmnimAbs.com

order@Omnimabs.com

506 N. GARFIELD AVE #210 ALHAMBRA, CA 91801

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt