

Catalog: OM109824



Scan to get more validated information

Rabbit anti-TAF15 polyclonal antibody - N-terminal region

Catalog: OM109824	
	100ug

Product profile

Product name Rabbit anti-TAF15 polyclonal antibody - N-terminal region

Antibody Type Primary Antibodies

Immunogen The immunogen for anti-TAF15 antibody: synthetic peptide directed towards the N terminal of human TA

F15

Key Feature

Clonality Polyclonal

Isotype IgG

Host Species Rabbit

Tested Applications WB

Species Reactivity Bovine Dog Guinea Pig Horse Human Mouse Pig Rabbit Rat

Concentration 1 mg/ml

Purification Affinity purified

Target Information

Gene Symbol TAF15

Gene Synonyms Npl3; RBP56; TAF2N; TAFII68; hTAFII68

Gene Full Name TAF15 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 68kDa

Gene Summary Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The p

rotein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. T AFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modi

fy general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. Its gene

encodes a subunit of TFIID present in a subset of TFIID complexes. Translocations involving chromosom e 17 and chromosome 9, where the gene for the nuclear receptor CSMF is located, result in a gene fusion product that is an RNA binding protein associated with a subset of extraskeletal myxoid chondrosarcoma s. Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promot er to position the polymerase properly, serves as the scaffold for assembly of the remainder of the trans cription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes a subunit of TFIID present in a subset of TFIID complexes. Translocations involving chromosome 17 and chromosome 9, where the gene for the nuclear receptor CSMF is located, result in a gene for usion product that is an RNA binding protein associated with a subset of extraskeletal myxoid chondrosa recomas. Two transcripts encoding different isoforms have been identified.

Alternative Names

Npl3, RBP56, TAF2N, TAFII68, hTAFII68

Molecular Weight (MW)

62kDa

Sequence

592 amino acids

Database Links

Entrez Gene 8148

SwissProt ID Q92804

Protein Accession NP 631961

Application

Western blot

0.2-1 ug/ml

ELISA Titer: 1:62500

Positive Control: HepG2 cell lysate

Application Notes WB:1:500~1:2000

 $\textbf{Notes}: Optimal\ dilutions/concentrations\ should\ be\ determined\ by\ the\ researcher.$

Additional Information

Form Liquid

Storage Instructions Aliquot and store at -20°C. Avoid repeated freeze / thaw cycles

Storage Buffer phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.

Note The product is for research use only,not for use in diagnostic or therapeutic procedures.

OmnimAbs.com

order@Omnimabs.com

506 N. GARFIELD AVE #210 ALHAMBRA, CA 91801

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt