



Catalog: OM161189

Scan to get more validated information



Syk Antibody (LR)

Catalog: OM161189

☐ 100ul

Product profile

Product name	Syk Antibody (LR)
Antibody Type	Primary Antibodies
Modification Notes	Syk (spleen tyrosine kinase) is a 635 amino acid protein that contains one protein kinase domain and two SH2 domains. One of several members of the protein kinase superfamily, Syk functions as a positive effector of B cell antigen receptor (CD79)-stimulated responses, coupling CD79 with the movement of one calcium ion through one of two phospho-regulated pathways. Specifically, calcium ions travel through either a phosphoinositide 3-kinase (PI 3-kinase)-dependent pathway when Syk is not phosphorylated, or through a phospholipase C (PLC) Ca^{2+} -dependent pathway when human Syk is phosphorylated on Tyr 348 and Tyr 352. Via its ability to influence CD79 activity and to control the movement of calcium through the cell, Syk plays an important role in a variety of cellular responses, including differentiation, phagocytosis, proliferation and B cell development. Syk expression is upregulated in T cell lymphoma, suggesting a possible role for Syk in tumorigenesis. Two isoforms of Syk, designated short and long, exist due to alternative splicing events.

Key Feature

Clonality	Polyclonal
Isotype	IgG
Host Species	Rabbit
Tested Applications	WB ,IP ,IF ,FC ,ELISA
Species Reactivity	Human Mouse Rat
Concentration	1mg/ml
Purification	Affinity purified


Target Information

Alternative Names	epitope mapping within the "linker" region of Syk of human origin
Tissue Specificity	epitope mapping within the "linker" region of Syk of human origin


Database Links

Entrez Gene 6850


Application




Application
Western blot analysis of Syk expression in non-transfected: (A) and human Syk transfected: (B) 293 whole cell lysates.



Application
Western blot analysis of Syk expression in non-transfected: (A) and human Syk transfected: (B) 293 whole cell lysates.



Application
Intracellular FCM analysis of fixed and permeabilized BJAB cells. Black line histogram represents the isotype control, normal rabbit IgG: .



Application
Western blot analysis of Syk expression in BJAB (A) and Ramos (B) whole cell lysates.

Additional Information

Form	Liquid
Storage Instructions	For short-term storage, store at 4° C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Storage Buffer	phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Note	The product is for research use only,not for use in diagnostic or therapeutic procedures.

OmnimAbs.com

order@Omnimabs.com
506 N. GARFIELD AVE #210 ALHAMBRA, CA 91801
This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt