Human BGH3 Recombinant protein

Catalog No.: OM770638

Sequence Information

Species: Human **Gene ID:**7045

Swiss Prot:Q15582 **Synonyms:** IgG3 TGFBI BIGH3 Beta ig-h3 Transforming growth factor-beta-induced protein ig-h3 CDB1 CDG2 CDGG1 CSD EBMD LCD1

Residues: Met194-His683

MYQNSNIQIHHYPNGIVTVNCARLLKADHHATNGVVHLIDKVISTITNNIQQIIEIEDTFETLRAAVAASGLNTMLE GNGQYTLLAPTNEAFEKIPSETLNRILGDPEALRDLLNNHILKSAMCAEA IVAGLSVETLEGTTLEVGCS GDMLTINGKAIISNKDILATNGVIHYIDELLIPDSAKTLFELAAESDVSTAIDLFRQAGLGNHLSGSERLTLLAPLNSVF KDGTPPIDAHTRNLLRNHIIKDQLASKYLYHGQTLETLGGKKLRVFVYRNSLCIENSCIAAHDKRGRYGT LFTMDRVLTPPMGTVMDVLKGDNRFSMLVA AIQSAGLTETLNREGVYTVFAPTNEAFRALPPRERSRLLG DAKELANILKYHIGDEILVSGGIGALVRLKSLQGDKLEVSLKNNVVSVNK EPVAEPDIMA TNGVVHVITN VLQPPANRPQERGDELADSALEIFKQASAFSRASQRSVRLAPVYQKLLERMKH

Product Information

Source: Prokaryotic expression.

Host: *E. coli*

Tags: N-terminal His-Tag

Subcellular Location: Secreted.

Purity: >95%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, 5% Trehalose.

Original Concentration: 200μg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.22

Predicted Molecular Mass: 53.6kDa

Accurate Molecular Mass: 60kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH_2O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious

degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]




Figure 1. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.